

Black Hole Zeckendorf Games

Caroline Cashman, Steven J. Miller, Jenna Shuffelton and Daeyoung Son

Abstract. Let F_n denote the n^{th} Fibonacci number (with first few terms 1, 2, 3, 5). The Zeckendorf game starts with N chips on pile F_1 , and on a turn a player may do one of two moves: if there are chips on piles F_k and F_{k-1} they may remove those and place one on F_{k+1} , or if there are two chips on F_k they may remove those and put one on F_{k+1} and one on F_{k-2} (unless $k \leq 2$: if there are two chips on F_2 we replace that with one each on F_3 and F_1 , while if there are two on F_1 we replace with one on F_2). Note at any moment in the game if c_n is the number of chips on pile F_n then $\sum c_n F_n = N$. Two players alternate turns until no moves remain. It was quickly proved that if $N > 2$ then Player 2 has a winning strategy and the game ends in the Zeckendorf decomposition of N (i.e., the representation of N as a sum of non-adjacent Fibonacci numbers); unfortunately it's a non-constructive proof. Subsequent papers proved many results about the game and generalizations, but the winning strategy remains a mystery. We thus consider a new variant: the Black Hole Zeckendorf game, where there is a fixed m such that if a chip is placed on pile F_m it becomes inaccessible (we also allow the two players to add chips to set up the initial board, so we do not have to start with N chips on F_1). We constructively find winning strategies for some m , where the winner depends on $N \bmod 16$.

Keywords. Zeckendorf decompositions, Fibonacci numbers, game theory, recreational mathematics.

2010 Mathematics Subject Classification. 91A46, 11P99.

1 The Zeckendorf Game

The beauty of the Fibonacci numbers is undeniable: a simple sequence, recursively defined by the sum of the two previous numbers, that has the tendency to show up in both natural and surprising places. Indexing so that $F_1 = 1$, $F_2 = 2$ and $F_{k+1} = F_k + F_{k-1}$, Zeckendorf proved a particularly interesting fact about the Fibonacci numbers, namely that any positive integer n can be written as the sum of non-adjacent Fibonacci numbers, known as the number's Zeckendorf decomposition [Ze72]. Baird-Smith, Epstein, Flint, and Miller [BEFM20, BEFM19], created a game from the process of converting a positive integer into its Zeckendorf decomposition using the moves of $F_i + F_{i-1} = F_{i+1}$ and $2F_i = F_{i+1} + F_{i-2}$, where F_i is the i^{th} Fibonacci number. We outline the rules to the original Zeckendorf game as follows.

1. **Setup:** The game is played on a board with columns corresponding to each of the Fibonacci numbers, indexing so that the 1st column corresponds with $F_1 = 1$, the 2nd column corresponds with $F_2 = 2$ and the m^{th} column corresponds with F_m , the m^{th} Fibonacci number. All n pieces begin in the 1st column.
2. **Gameplay:** Players alternate, selecting their moves from the following.
 - (a) Adding consecutive terms: If the board contains pieces in both F_i and F_{i-1} columns, players can remove one piece from each column to add as one piece in the F_{i+1} column.
 - (b) Merging 1's: If the board contains more than one piece in the F_1 column, players can remove two pieces from the F_1 column to merge as one piece in the F_2 column.
 - (c) Splitting: If the board contains more than one piece in the F_2 column, players can split two pieces from the F_2 column to place one piece in each F_1 and F_3 . For $i \geq 3$, players can split two pieces in the F_i column to place one in each F_{i-2} and F_{i+1} .
3. **Winning:** The last player to move wins.





Figure 1: Game tree for $n = 14$, with a winning path in green, as included in [BEFM20].

They proved that the game is playable, meaning it always ends in finite time, and that the final board placed down will be equal to the Zeckendorf decomposition of n . Moreover, they showed that for all $n \neq 2$, Player 2 has a winning strategy. Notably, this is not a constructive winning strategy, and instead relies on a parity stealing argument. If one assumes that Player 1 has a winning strategy, Player 2 later has the opportunity to steal it, therefore Player 2 must have a winning strategy. With increasing n , the number of possible game positions grows exponentially, making the construction of a winning solution for Player 2 challenging. Even for n as small as 14, as in Figure 1, it is not obvious how Player 2 wins without use of brute force.

2 The Black Hole Zeckendorf Game

In order to better understand the original Zeckendorf game, we consider a variation of the game occurring on a smaller board. We call this the “ F_m Black Hole” variation of the Zeckendorf game, where once a piece is placed on some F_i for $i \geq m$, it falls into the “Zeckendorf Black Hole” and is permanently removed from game play. This variant reduces the number of possible moves a player has, making the game easier to analyze. Here, a game played with n pieces ends in the Zeckendorf decomposition of $n \pmod{F_m}$. As we play through games, we let bold blue denote Player 2 and red denote Player 1. We focus on the case with a black hole on $F_4 = 5$, which allows for the following moves:



We also consider an empty board phase of the game in which players take turns placing pieces

	$a \equiv 0 \pmod{3}$	$a \equiv 1 \pmod{3}$	$a \equiv 2 \pmod{3}$
$c \equiv 0 \pmod{4}$	$\alpha \geq \gamma$ $\alpha \leq \gamma - 1$	$\forall \alpha, \gamma$	$\alpha \geq \gamma + 1$ $\alpha \leq \gamma$
$c \equiv 1 \pmod{4}$	$\alpha \geq \gamma - 1$ $\alpha \leq \gamma - 2$	$\forall \alpha, \gamma$	$\alpha \geq \gamma$ $\alpha \leq \gamma - 1$
$c \equiv 2 \pmod{4}$	$\forall \alpha, \gamma$	$\alpha \geq \gamma + 1$ $\alpha \leq \gamma$	$\forall \alpha, \gamma$
$c \equiv 3 \pmod{4}$	$\forall \alpha, \gamma$	$\alpha \geq \gamma$ $\alpha \leq \gamma - 1$	$\forall \alpha, \gamma$

Figure 2: Winners for board setups $(a, 0, c) = (3\alpha + k_1, 0, 4\gamma + k_3)$ with $\alpha, \gamma, k_1, k_3 \in \mathbb{Z}^{\geq 0}$, $0 \leq k_1 \leq 2, 0 \leq k_3 \leq 3$ in an F_4 Black Hole Zeckendorf Game. Player 2 wins are depicted in bold blue, and Player 1 wins are depicted in red.

in the outermost columns, with the last player to place assuming the role of Player 2 during the decomposition phase of the game. Defining this phase of the game as such allows players to use move mirroring as a strategy. In order to force an advantageous set-up, players can place pieces in the column opposite their opponent. Combining these two variations of the original game, we find the following main result.

Theorem 21. *Player 2 has a constructive strategy for winning an Empty Board F_4 Black Hole Zeckendorf game for any $n \equiv 0, 2, 4, 6, 9, 11, 13 \pmod{16}$ such that $n \neq 2, 32$ in which case Player 1 has the winning strategy. Player 1 has a constructive strategy for winning an Empty Board F_4 Black Hole Zeckendorf game for any $n \equiv 1, 3, 5, 7, 8, 10, 12, 14, 15 \pmod{16}$, such that $n \neq 17, 47$, in which case Player 2 has the winning strategy.*


To provide a constructive solution, we determine winning positions for a board setup $(a, 0, c)$ as outlined in Figure 2, and then provide a path from winning position to winning position. Knowing which player wins for a given setup allows us to determine which player move mirroring benefits, giving us our winners as stated in Theorem 21. However, it is not immediately clear that the positions in the table are in fact winning positions. In order to show that they are, we show that certain players win key intermediate positions using nonconstructive methods. See, for example Lemma 22, which uses the following corollary:

Corollary 21. *Let (a, b, c) be a game state for an F_4 Black Hole Zeckendorf game. Player 2 has a winning strategy at $(1, 0, c)$ for all $c \neq 3 \in \mathbb{Z}^{\geq 0}$ and at $(0, 1, c)$ for all $c \neq 1, 2, 6 \in \mathbb{Z}^{\geq 0}$. Player 1 has a winning strategy at $(2, 0, c)$ for all $c \neq 1 \in \mathbb{Z}^{\geq 0}$ and at $(2, 1, c)$ for all $c \in \mathbb{Z}^{\geq 0}$.*

Lemma 22. *Let (a, b, c) be a game state for an F_4 Black Hole Zeckendorf game. For all $\alpha, \gamma, k_1, k_3 \in \mathbb{Z}^{\geq 0}$, such that $1 \leq k_1 \leq 2$, and $0 \leq k_3 \leq 3$, Player 1 has a winning strategy for $(3\alpha + k_1, 1, 4\gamma + k_3)$.*

Proof. We proceed by a non-constructive proof. For contradiction's sake, suppose Player 2 wins $(3\alpha + k_1, 1, 4\gamma + k_3)$ when $k_1 = 1, 2$. Then, consider the game tree in Figure 3, where r is the number of rounds in which Player 1 then the Player 2 plays. We note that Player 1 has other possible moves, but we only consider moves relevant to the proof.

By assumption, Player 2 has a winning strategy regardless of what the other player places. Suppose Player 1 places $(3\alpha + k_1 - 1, 0, 4\gamma + 1)$ for their first move. Then, Player 2 can place

Figure 3: Game Tree for the Game State $(3\alpha + k_1, 1, 4\alpha + k_3)$.

either $(3\alpha + k_1 - 3, 1, 4\gamma + 1)$ or $(3\alpha + k_1, 0, 4\gamma - 1)$. But as shown in the tree, Player 1 had the opportunity to place $(3\alpha + k_1, 0, 4\gamma - 1)$ in the round before; so by assumption, placing it is a losing move. It follows that in order to win, Player 2 must place $(3\alpha + k_1 - 3, 1, 4\gamma + 1)$. As shown in the tree, Player 1 has the same options as before, so if they place $(3\alpha + k_1 - 4, 0, 4\gamma + k_3 + 2)$, then by assumption, Player 2 must place $(3\alpha + k_1 - 6, 1, 4\gamma + k_3 + 2)$.

Then, the game eventually reduces down to Player 2 placing $(3\alpha + k_1 - 3r, 1, 4\gamma + k_3 + r)$ after r rounds of Player 1 then Player 2 placing. After the α^{th} round, Player 2 will place $(k_1, 1, 4\gamma + k_3 + \alpha)$. If $k_1 = 1$, then Player 1 can add from columns F_2 and F_3 to place $(1, 0, 4\gamma + \alpha + k_3 - 1)$ which wins by Corollary 21 for all $4\gamma + \alpha + k_3 - 1 \neq 3$. If $k_1 = 2$, then Player 1 can add from columns F_1 and F_2 to place $(1, 0, 4\gamma + \alpha + k_3 + 1)$ which wins by Corollary 21 for all $4\gamma + \alpha + k_3 + 1 \neq 3$. Additionally, we show in the Appendix in [CMSS24], that Player 1 also has a winning strategy for the cases when $4\gamma + \alpha + k_3 \pm 1 = 3$. This is a contradiction to the assumption that Player 2 has a winning strategy, so then Player 1 must have some winning strategy when $(3\alpha + k_1, 1, 4\gamma + k_3)$ is placed for $k_1 = 1, 2$.

Through similar proofs that show which players win intermediate positions, we are able to show that Figure 2 accurately describes which player wins a given position. From here, we provide a path from winning position to position, thereby providing a constructive solution to the game.

3 Example Winning Strategy

Let us reconsider the case when $n = 14$, but this time when playing the Empty Board Black Hole Zeckendorf Game with a Black Hole on F_4 . If Player 1 places their first piece in the first column, and then mirrors Player 2's moves, then they can force Player 2 to set the board down as $(5, 0, 3)$. Player 1 can then use the strategy below to limit Player 2's moves, therefore leading to a Player 1 win.

$$\begin{array}{c}
 (5, 0, 3) \\
 | S_3 \\
 (6, 0, 1) \\
 | M \\
 (4, 1, 1) \\
 | A_2 \\
 (4, 0, 0) \\
 | M \\
 (2, 1, 0) \\
 | A_1 \\
 (1, 0, 1)
 \end{array}$$

4 Future Work

For our definition of the empty board game, it was only necessary to analyze which player won for board setups $(a, 0, c)$. Allowing players to place pieces in the center column during the empty board phase could be an interesting area for future work, as the strategy of move mirroring is no longer as applicable and there are more possible setups for the decomposition phase of the game to analyze.

Note also that for any $m \geq 5$, the strategy of reducing the game modulo F_m is not as immediately successful, as it is challenging for either player to place a piece in the black hole without first giving another player the opportunity to do so.

Acknowledgement. Authors were supported by Williams College, The Finnerty Fund, The College of William & Mary Charles Center, and NSF Grant DMS2241623. Special thanks to Paul Baird-Smith, whose code we edited to play through the Zeckendorf Black Hole game.

References

- [BEFM20] P. Baird-Smith, A. Epstein, K. Flint and S. J. Miller, *The Zeckendorf Game*, Combinatorial and Additive Number Theory III, CANT, New York, USA, 2017 and 2018, Springer Proceedings in Mathematics & Statistics **297** (2020), 25–38.
- [BEFM19] P. Baird-Smith, A. Epstein, K. Flint and S. J. Miller, *The Generalized Zeckendorf Game*, Proceedings of the 18th International Conference on Fibonacci Numbers and Their Applications, Fibonacci Quarterly **57** (2019), no. 5, 1–14
- [CMSS24] C. Cashman, S. J. Miller, J. Shuffelton, D. Son, *Black Hole Zeckendorf Games*, preprint 2024 (to appear in the Fibonacci Quarterly).
- [Ze72] E. Zeckendorf, *Représentation des nombres naturels par une somme des nombres de Fibonacci ou de nombres de Lucas*, Bulletin de la Société Royale des Sciences de Liège **41** (1972), 179–182.

Caroline Cashman

Department of Mathematics
College of William & Mary, Williamsburg, VA

e-mail: cecashman@wm.edu

Steven J. Miller

Department of Mathematics and Statistics,
Williams College, Williamstown, MA

e-mail: sjm1@williams.edu

Jenna Shuffelton

Department of Mathematics and Statistics,
Williams College, Williamstown, MA

e-mail: jms13@williams.edu

and **Daeyoung Son**

Department of Mathematics and Statistics,
Williams College, Williamstown, MA

e-mail: ds15@williams.edu